

See discussions, stats, and author profiles for this publication at: <https://www.researchgate.net/publication/316188170>

Developmental abnormalities of *Betta splendens* embryos reared in oil and grease contaminated water samples of Vavuniya reservoir, Sri Lanka

Article · April 2017

CITATIONS

0

READS

101

4 authors:



Umasuthan Srivignesvaran

University of Peradeniya

3 PUBLICATIONS 0 CITATIONS

SEE PROFILE



Anton Emmanuel Selvakumar Patrick

University of Jaffna

36 PUBLICATIONS 26 CITATIONS

SEE PROFILE



Naveendrakumar Gnanapragasam

University of Jaffna

15 PUBLICATIONS 55 CITATIONS

SEE PROFILE



Sivashanthini Kuganathan

University of Jaffna

81 PUBLICATIONS 252 CITATIONS

SEE PROFILE

Some of the authors of this publication are also working on these related projects:



population dynamics of *Gerres oblongus* [View project](#)



biology of scomberoides [View project](#)



Developmental abnormalities of *Bettasplendens* embryos reared in oil and grease contaminated water samples of Vavuniya reservoir, Sri Lanka

Umasuthan S.^{1*}, Patrick A.E.S.¹, Naveendrakumar G.¹ and Kuganathan S.²

¹Department of Bio-Science, Faculty of Applied Science, Vavuniya Campus of the University of Jaffna, Sri Lanka

²Department of Fisheries, Faculty of Science, University of Jaffna, Sri Lanka
umasuthan.sm@gmail.com

Available online at: www.isca.in, www.isca.me

Received 27th February 2017, revised 3rd April 2017, accepted 9th April 2017

Abstract

In Vavuniya urban area the wash-outs from petroleum automobile service stations contain Kerosene oil, lubricating engine oil, grease as major compounds that contributing oil and grease pollution in Vavuniya reservoir, Sri Lanka. To investigate the potential oil and grease toxicity on fish embryo development, the *Bettasplendens* (Siamese fighting fish) embryos were reared in water samples from Vavuniya reservoir. Average value of oil and grease content from polluted location was determined for the preparation of artificial aliquots of oil and grease (20 mg of Engine oil: Kerosene oil: Grease = 1:1:1 w/w). These aliquots mixed with the 1000 ml of water samples obtained from non-polluted location of this reservoir. Prepared medium was shaken vigorously for 5 minutes and 25ml was added in each Petridish to prepare the larval growth medium. Meantime, 25ml of non-polluted water was taken in another set of Petri dishes as a control setup. Treatments and control were allowed under natural diffuse photo periods. Frequent microscopic examinations were done until 72hpf (hours post-fertilization) for embryological abnormalities and compared with control group from March to December 2016. Significant embryo developmental abnormalities such as delayed and reduced outgrowth of the caudal fin-fold ($p = .008$), reduction in melanin pigment formation ($p = .04$) and higher larval mortality (75 %) were observed. Hence, much higher potential of oil and grease impacts on fish embryo development observed in this study, suggests the threats for survival of local fish species in Vavuniya reservoir. Assessment of oil and grease impact on the reproductive potential of the local fish species is necessary for its conservation in their natural habitats.

Keywords: Developmental abnormalities, Oil and grease pollution.

Introduction

Most of the studies on phototoxicity of polycyclic aromatic compounds (PAC) in water bodies have done on zooplanktons¹. However, only a few researches have focused on phototoxicity of PAC preparations in fish embryo development and among this, majority of research works examining the mechanism of this toxicity in fish early life stages, especially on undefined crude oil². Though, minority of studies have only addressed the oil and grease toxicity³. Tricyclic fluorenes, dibenzothiophenes and phenanthrenes are the most abundant PAC that acts as cardiotoxicagents and causes pericardial and yolk sac edema on fish larvae^{5,6}. Although, researchers found some toxic effects like intracranial bleeding and blister-like abnormalities of the caudal fin tissues in Zebra fish embryos resulting from oil exposure, the exact pathological origin is unknown⁶. These oil and grease (O&G) compounds are poly-aromatic hydrocarbons (PAHs) and containing toxic impurities which could cause the harmful effects to the aquatic organisms, such as in Vavuniya reservoir, Sri Lanka. As this reservoir is situated in the urban area, experiencing frequent oil and grease pollution that caused by petroleum automobile service stations. Kerosene like odour in *Oreochromis niloticus* due to oil and grease pollution in Vavuniya reservoir causes negative impacts on inland fisheries

in recent years⁴. In order to study the effects of O&G on fish embryo development and larval growth performance, Siamese fighting fish (*Bettasplendens*) was selected. Indeed, *Bettasplendens* have showed a viable growth performance and higher percentage of survival in the laboratorial conditions in the northern region of Sri Lanka⁷. Hence, *Bettasplendens* is an ideal oviparaous fish species with aerial respiratory adaptations for laboratory oriented studies.

Materials and methods

Sample collections: Water samples were collected fortnightly from polluted location of the Vavuniya reservoir using water sampler (grab sampling methods was adapted) and average content of oil and grease was determined by using gravimetric method.

Determination of oil and grease: The samples collected from the polluted location of the Vavuniya reservoir were extracted with three aliquotes of Hexane in a separator funnel. Samples were shaken vigorously for 2 minutes per extraction. The first Hexane aliquot was used to rinse the sample container so that its entire contents were transferred to the extraction vessel. The ratio of solvent to sample was 1:10, i.e. 20 ml of Hexane (per

extraction) per 200ml of sample. The solvent extracts were passed through a drying funnel containing anhydrous sodium sulfate to absorb the additional moisture. Then the oil and grease combined with Hexane was separated using rotary evaporator then the weight of the oil and grease was determined after being kept at desiccators overnight.

Artificial preparation of oil and grease: According to the average value of O&G content of Vavuniya reservoir water sample (polluted location), artificial aliquots of O&G were prepared (set to resemble the sample water from Vavuniya reservoir polluted location in terms of weight in parts per million). Artificially prepared oil and grease of 20mg (Engine oil: Kerosene oil: Grease=1:1:1 w/w) was mixed with the 1000 ml of water samples obtained from non-polluted location of this reservoir where, embryos to be reared as treatment. The prepared water medium was shaken vigorously for 5 minutes and 25ml was placed in each Petridish to prepare the larval growth medium. As a control setup, 25ml of non-polluted water (without O&G contaminations) was taken in another set of Petri dishes. Three replicates were made for treatments and control and allowed under natural diffuse photo periods. Frequent microscopic examinations were done until 72 hpf (hours post-fertilization) for embryological abnormalities and analyzed with control group from March to December 2016.

Bettasplendens embryo exposure: Onset of spawning of *Bettasplendens*, embryos were transferred to the Petridishes at a rate of 10 embryos per Petridish by using delicate droppers. Mild aeration was initiated which resembles the semi-artificial incubators at room temperature under natural diffuse photo

periods (Figure-1). Assumptions were made that all the eggs were fertilized and viable onset of spawning.

Microscopic imaging: Frequent microscopic observations were done until 72hpf (hours post-fertilization) from early embryos (zygote) via fry to early larval stages³. Abnormalities among tail morphometry (caudal fin outgrowth) and pigmentation intensity were analyzed by using XENON Student's Microscopes. Microscopic images of embryo development from control and experimental setup were imaged using digital camera (Sony DSC-W350 14.1 Mega Pixel, 72dpi). Mortality rates were recorded by direct observations from 6 – 120 hpf.

Statistical analysis: To compare the morphometric differences in control and in treatment, 2t –Proportion test was performed for the ratios of Normal: Reduced fin-folds in both control and treatment. This analysis was performed (at 95% confident interval, $\alpha=0.05$) using Minitab 16.0) to find-out the significant differences.

Results and discussion

The fish embryo developmental abnormalities such as delayed and reduced outgrowth of the caudal fin-fold, reduction in melanin pigment formation were significantly ($p=.04$) greater in the oil and grease contaminated water sample (20 ppm) than the control. Onset of fertilization, active cell division started in the blasto-disc of fish eggs (Figure-2: B₁, B₂) in all setups. Though cytolysis occurred in the oil and grease contaminated water sample (treatment) (Figure-2: A₁, A₂). Percentage of larval mortality was higher in the treatment than control.




Figure-1: Artificially prepared incubator units for fish embryo development. [Control setup with non-polluted water samples, Treatment with O&G constituents]




Figure-2: Dead eggs were opaque (A₁, A₂). Viable eggs are clear with prominent previtteline space and active cell division visible in the blasto-disc in the animal pole (B₁, B₂).

The embryo development in the control setup was found as the ratio of normal fin-fold: reduced fin-fold growth is 9:1 and the percentage of larval mortality is 20% (Table-1). This reduced fin-fold growth phenomena of few individuals may be due to the genetic influences (in-born). However, majority of the embryos showed viable growth performance in the experimental setup.

In treatment, higher larval mortalities (75%), reduced fin-fold ratio and less pigmentation were observed. This was suggested, mainly due to the acute toxicity of oil and grease constituents. There was a significantly strong difference (p = .008) observed in normal: reduced caudal fin-folds (Figure-3) between control and treatment.

Pigment intensity (Figure-3) was also higher in control than treatment in this analysis (Table-2).

There was not any remarkable pericardial edema, yolk sac edema and intra cranial hemorrhaged observed in *Bettasplendens* (Figure-4).

Table-1: Summary of embryo development of fish (*Bettasplendens*).

Developmental variables	Control	Treatment
Number of embryos	120	120
Survival rate (%)	80	25
Mortality rate (%)	20	75
Normal fin-fold	86	20
Reduced fin-fold	10	10
Ratio; Normal: Reduced	9:1	2:1
High pigmentation	84	21
Low pigmentation	12	9
Ratio : high: low	7:1	7:3

Table-2: Summary of statistical analysis of 2t-Propotion test for caudal fin & pigmentation.

Factors	Treatment (Observed values)	Control (Expected values)	p-values
Caudal fin:			0.008
Normal outgrowth	13	58	
Reduced outgrowth	07	06	
Melanin pigmentation:			0.042
High pigment intensity	21	84	
Low pigment intensity	09	12	

Comparatively lesser melanin pigment formation and reduced caudal fin outgrowth of *Bettasplendens* larvae was clearly visible in the oil and grease contaminated water sample than the control (Figure-3 and 4). This kind of effects observed in the Zebra fish larvae kept in crude oil aliquots with natural sunlight exposure³.

Poly-aromatic hydrocarbons (PAHs) and toxic impurities from petroleum automobile service stations caused harmful effects to the fish early life histories. Indeed, the embryo development of *Bettasplendens* highly affected by these compounds under indoor laboratorial conditions with diffuse natural photo periods. Especially, embryo development of the fish is highly affected by these compounds⁶. Reduced caudal fin outgrowth of Zebra fish embryos reared in the crude oil treatment in the presence of solar radiation has been proved in other studies³.




Figure-3: Morphological abnormalities of *Bettasplendens* larvae (x 100) in control (1) and treatment (2): Variation in pigment intensity (dorsal view), higher pigmentation in A₁ and B₁ comparing with A₂ and B₂ (white arrows indicates the location of pigmentations). C & D: Morphology of caudal-fin outgrowth (lateral view); higher caudal-fin outgrowth in C₁ and D₁ comparing with C₂ and D₂ (black arrows indicates the outline of the caudal-fin).



Figure-4: Newly hatched fry (x 100) of *Bettasplendens* in the treatment set up where the caudal fin out growth is reduced but pericardial and yolk sac edema were not clearly visible.

Conclusion

Reduced caudal fin outgrowth, reduction in melanin pigment formation and higher larval mortality rates justifies the acute toxicity of oil and grease on *Bettasplendens* embryo development. Hence, much higher phototoxic potential of oil and grease impacts on fish embryo development observed in this study, suggests the threats for survival of local fish species in Vavuniya reservoir. Thus, understanding the consequences of the consequences of varying oil and grease discharges from petroleum automobile service stations on abnormalities of local fish species in inland water bodies are necessary for safeguard the rich fish species composition and abundance.

References

1. Cleveland L., Little E., Calfee R.D. and Barron M.G. (2000). Photoenhanced toxicity of weathered oil to *Mysidopsis bahia*. *Aquat. Toxicol.*, 49(1), 63-76.
2. Barron M.G., Carls M.G., Short J.W. and Rice S.D. (2003). Photoenhanced toxicity of aqueous phase and chemically dispersed weathered Alaska North Slope crude oil to Pacific herring eggs and larvae. *Environ. Toxicol. Chem.*, 22(3), 650-660.
3. Hatlen K., Sloan C.A., Burrows D.G., Collier T.K., Scholz N.L. and Incardona J.P. (2010). Natural sunlight and residual fuel oils are an acutely lethal combination for fish embryos. *Journal of Aquatic Toxicology*, 99(1), 56-64.
4. Patrick A.E.S., Kuganathan S. and Edirisinghe U. (2014). Linear morphometric phenomenon of *Oreochromis niloticus* in polluted location of Vavuniya reservoir, Sri Lanka. Proceedings of the Postgraduate Institute of Science Research Congress, University of Peradeniya, Sri Lanka. 84.
5. Carls Mark G., Holland Larry, Larsen Marie, Collier Tracy K., Scholz Nathaniel L. and Incardona John P. (2008). Fish embryos are damaged by dissolved PAHs not oil particles. *Aquatic Toxicol.*, 88(2), 121-127.
6. Incardona J.P., Carls M.G., Teraoka H., Sloan C.A., Collier T.K. and Scholz N.L. (2005). Aryl hydrocarbon receptor-independent toxicity of weathered crude oil during fish development. *Environ. Health Perspect.*, 113(12), 1755-1762.
7. Patrick A.E.S. (2009). Viable growth performance of *Bettasplendens* (Regan, 1910) larvae with different aquaria environment and diets in Jaffna District. Sri Lankan Association for the Advancement of Science (SLAAS) on 65th Annual Session, 82.
8. Arfsten D.P., Schaeffer D.J. and Mulveny D.C. (1996). The effects of near ultraviolet radiation on the toxic effects of polycyclic aromatic hydrocarbons in animals and plants: a review. *Ecotoxicol., Environ. Saf.*, 33(1), 1-24.
9. Yu H. (2002). Environmental carcinogenic polycyclic aromatic hydrocarbons: photochemistry and phototoxicity. *Journal of Environmental Science Health C: Environ. Carcinog. Ecotoxicol. Rev.*, 20(2), 149-183.